arXiv:1209.0883v1 [hep-th] 5 Sep 2012

Heavy Quark Potential from Gauge/Gravity Duality: A Large D Analysis

Vikram Vyas[*]

Department of Physics, Shiv Nadar University,
Gautam Budh Nagar, 203207, UP, India.
Email: *visquare@gmail.com*

Abstract

Heavy quark potential is calculated in the framework of Gauge/Gravity duality using the large D approximation, where D is the number of dimensions transverse to the flux-tube connecting a quark and an anti-quark in a flat D+2 dimensional spacetime. We find that in the large D limit the leading correction to the ground state energy, as given by an effective Nambu-Goto string, arises not from the heavy modes but from the behavior of the massless modes in the vicinity of the quark and the anti-quark. This correction modifies the Arvis's formula for the ground state energy of a Nambu-Goto string by an overall factor which has a very mild dependance on the distance between the quark and the anti-quark. This analysis suggests that the leading corrections to an effective Nambu-Goto string description of a QCD flux-tube are in the form of boundary terms.

---

[*] On leave of absence from St. Stephen's College, Delhi, India

Contents

1 Introduction . . . 2
2 Heavy Quark Potential from Confining Geometry in the Large D Limit . . . 5
3 The $1/D$ Corrections from Heavy Modes . . . 12
4 The Boundary Effects . . . 14
5 Summary and Conclusions . . . 18
A Appendix: Fluctuations Around the Minimal Surface . . . 20

1 Introduction

There is strong numerical evidence that the dynamics of a flux-tube formed between a heavy quark and an anti-quark is well described by the Nambu-Goto string in four space-time dimensions (see for e.g [1, 2, 3].) In particular the ground state energy of such a flux-tube matches very well with the Arvis's formula for the ground state energy of an open Nambu-Goto string with fixed end-points in four dimensions [4]. Yet, one expects on general grounds that there should be corrections to Arvis's formula, for after all, the QCD flux-tube is not a one-dimensional object, it has some "intrinsic" thickness which we expect to be related to the *finite* correlation length of the confining $SU(3)$ gauge theory [5]. If one thinks of the fluctuations in the intrinsic thickness as an additional degree of freedom, then integrating out this degree of freedom would lead to an effective string action. Such an action, in general, should contain all possible terms consistent with reparameterization invariance. In the sprit of effective field theories, these terms can be organized into relevant, marginal and irrelevant terms (for a review of these ideas see, for e.g, [6, 7].) The Nambu-Goto action is the only relevant term, then there is a marginal term, the so called extrinsic-curvature term, and infinitely many irrelevant terms corresponding to higher and higher derivatives of the string coordinates. Further, since we are considering open strings, there could also be boundary terms. In fact we know on very general grounds that the resulting effective string action cannot be just the Nambu-Goto action [8].

To obtain an effective string action for QCD flux-tubes, starting from the fundamental description in terms of quarks and gluons, requires that we integrate out all the fluctuations of the gauge fields whose wavelength is less than the correlation length of the confining theory. A task which is as difficult as to obtain an effective chiral Lagrangian for pions starting from the QCD action. The gauge/gravity duality [9], inspired by the AdS/CFT correspondence [10, 11, 12, 13], provides us with an alternate, though approximate and often heuristic, way of analyzing the dynamics of strongly coupled gauge theories. Specifically, it provides us with a geometrical way of exploring the consequences of one of the defining properties of SU(N) gauge theories, namely the existence of a finite correlation length of the gauge fields. According to the gauge/gravity duality, the QCD flux-tube should be thought of as a holographic projection of the fundamental string in five-dimensional curved space. This picture then suggests that for the confining gauge theories the intrinsic thickness of the flux-tube can be related to the position of a fundamental string in the fifth dimension [14, 15]. It was also pointed out in [16] that this relationship was consistent with the expected behavior of the intrinsic thickness a Nielsen-Olesen like flux-tube with its string tension [17].

The aim of the present investigation is to use the geometrical picture provided by the gauge/gravity duality to delineate the various sources of correction to the heavy quark potential as compared to the ground state energy of a Nambu-Goto string in flat four dimensional space-time. These corrections are induced by the fact that the fundamental QCD string lives in a five-dimensional curved space. We will analyze this situation using the large D approximation, where $D+2$ is the number of dimensions of the flat spacetime where the quark lives. D is therefore the number of directions transverse to a QCD flux-tube. The main advantage that this approach offers is that it treats the massless modes of the string non-perturbatively.

In gauge/gravity duality, the confining property of the boundary gauge theory is reflected in the fact that a fundamental string in five dimension can minimize its energy by staying at a fixed value of its fifth coordinate, $Y = Y^*$, the coordinate which is transverse to the four dimensional flat Euclidian space, as shown in Fig.(1). But since the quark and the anti-quark live at the boundary, the end point of the fundamental string have to dip from its fixed value of $Y = Y^*$ to $Y = 0$, which is the boundary. We first calculate the large D behavior of the fundamental string in the approximation where the fundamental string dips from its minimal energy value to the boundary right at $x = 0$ and $x = L$, where the quark and the anti-quark are placed. This is shown in Fig.(2). The dynamics of the open string in this approximation is very much like that of an open string placed at $Y = Y^*$ and stretched between two d-branes. The leading term in the large D analysis for this case turns out to be nothing but the Arvis's formula. As we will see in section (4), when one corrects for this approximation, and takes into account that the fundamental string dips not right at $x = 0$ or $x = L$ but there is a finite transition region marked d in Fig. (1) then the leading term in the large D expansion is no longer the Arvis's formula alone. Rather the Arvis's formula gets modified by an overall factor which can be thought of as a length dependent string tension.

It is important to note there there should be corrections to the Arvis's formula even if the

Nambu-Goto action by itself provided an exact description of QCD flux-tubes [7, 18][1]. These expected corrections should arise from quantizing the Nambu-Goto string in non-critical dimensions. From the point of the large D expansion such corrections should exist since the Arvis's formula is only the leading term in the large D expansion and there is no a-prior reason to believe that the higher order corrections should vanish. Though we will not see such correction in our large D analysis, but it is interesting to note that these higher order corrections should themselves be proportional to $D - 24$ and therefore vanish for the critical dimension of $D + 2 = 26$. Thus, surprisingly the Arvis's formula gives the exact ground state energy of a Nambu-Goto string in flat space-time for $D = 24$ and for $D \to \infty$.

In an alternative approach, pioneered in [19, 20], one starts with a derivative expansion for the action of a fundamental string stretched between two d-branes in five dimensional confining background, then one perturbatively integrates out the fluctuations corresponding to the oscillations in the fifth dimension. These world-sheet fluctuations are massive, and after integrating them out one obtains an effective action for the massless transverse fluctuations. In [20] it was shown that this leads to a correction to the Arvis's potential at the order of $1/L^4$, where L is the length of the fundamental string stretched between the two d-branes. This correction comes from a boundary term induced by integrating out the heavy modes. In our large D analysis we do not see such a boundary term at least to the order of $1/D$, but its worth recalling that for the physical case of interest $D = 2$ and therefore even $1/D^2$ correction can be important. We do see boundary terms but their physical origin is very different. They come not from integrating out heavy modes but from the behavior of the massless modes near the boundary of the five-dimensional curved space.

The outline of the paper is as follows: In section 2, we consider a particular class of confining geometries and calculate the leading term in the large D expansion under the simplifying assumption that we discussed above. In section 3, within the same simplifying assumption, we calculate the $1/D$ correction arising from the fluctuation of the string in the fifth dimension which give rise to massive world-sheet modes. In section 4, we make allowance for the fact that the fundamental string reaches the boundary not suddenly at $x = 0$ and $x = L$ but smoothly over a transition region of finite length, d, which is a function of the distance L between the quark and the anti-quark. This leads to an effective string description of the QCD flux-tube in which the Nambu-Goto string action is supplemented by boundary terms. We summarize and state our conclusions in the final section.

---

[1] I would like to thank Ofer Aharony for pointing this out to me.

2 Heavy Quark Potential from Confining Geometry in the Large D Limit

The gauge invariant observable of a SU(N) gauge theory that is directly related to the heavy quark potential is the Wilson loop,

$$W\left[\Gamma\right] = \mathrm{Tr}\hat{\mathrm{P}}\left(\exp i\int_{\Gamma} A\right). \tag{2.1}$$

Here Γ is a closed curve in four dimensional Euclidean space, and A is the vector potentials of SU(N) gauge theory [21]. According to the assumed Gauge/Gravity duality, the expectation value of the Wilson loop can be written as a sum over the world-sheets of an open string whose end points terminate on the loop Γ. The open string itself lives in a five dimensional curved space whose boundary is the the four dimensional Euclidian space where Γ is located [13, 22]. Formally,

$$< W[\Gamma] >_{YM} \equiv \int [dA] \exp\left\{-S[A]_{YM}\right\} = \int_{\partial X=\Gamma} [dX] \exp\left\{-S[X]_{NG}\right\}, \tag{2.2}$$

here $S[A]_{YM}$ is the Yang-Mills action, while $X(\tau,\sigma)$ represents the world sheet of a string in a curved five dimensional space and $S[X]_{NG}$ is the Nambu-Goto action. To evaluate the Nambu-Goto action for a given surface in this space we need to know its metric. Ideally we would like a prescription which gives us the geometry of the five-dimensional space once the beta function (governing the running of the Yang-Mills coupling constant with energy) of the dual boundary quantum field theory is specified. In the absence of such a prescription we can look for simple modifications of AdS_5 space that leads to an area law for the expectation value of the Wilson loop. Following [15] we will consider one such class of modification, where the metric of the curved five dimensional space in the coordinate system $\{t, x, \mathbf{X}, Y\}$ is given by

$$ds^2 = F(Y)\left(dt^2 + dx^2 + d\mathbf{X}^2 + dY^2\right). \tag{2.3}$$

In this coordinate system $Y = 0$ is the four dimensional boundary[2] where the gauge theory lives and $\mathbf{X}$ denotes $D = 2$ directions transverse to the flux-tube lying along the x axis, while t is the Euclidian time. As, mentioned in the introduction, the area law for the expectation value of the Wilson loop is implemented by requiring that $F(Y)$ has a minima at some fixed value of $Y = Y^*$. For this class of metric we will evaluate (2.2) using a large D expansion, where D is the number of dimensions transverse to the QCD flux-tube.

---

[2] Strictly speaking for AdS_5 like space, $Y = 0$ is a conformal boundary [12]. $F(Y)$ diverges there, so in what follows we will assume that the boundary is at $Y = \epsilon$ and that corresponds to considering the boundary gauge-theory with a UV cutoff of the order $1/\epsilon$ [23]

In the static gauge, or a physical gauge, a world-sheet of the string connecting the quark and the anti-quark is given by

$$X = (t, x, \mathbf{X}(t,x), Y(t,x)) = (\tau, \sigma, \mathbf{X}(\vec{\sigma}), Y(\vec{\sigma})), \tag{2.4}$$

$$\mathbf{X} = \left(X^1, X^2, \ldots, X^D\right), \tag{2.5}$$

where we identify world sheet parameter τ with t and σ with x. A point on the world-sheet with coordinates (τ, σ) will be represented by $\vec{\sigma}$. The world-sheet coordinate range from

$$-\frac{T}{2} \leq t = \tau \leq \frac{T}{2}; \quad 0 \leq x = \sigma \leq L, \tag{2.6}$$

corresponding to a Wilson loop made up of the worldlines of a quark at $x = 0$ and an antiquark at $x = L$, and we will be interested in the limit $T \to \infty$. Surfaces appearing in (2.2) satisfy the boundary conditions:

$$\mathbf{X}(\tau, 0) = 0 = \mathbf{X}(\tau, L); \quad Y(\tau, 0) = 0 = Y(\tau, L). \tag{2.7}$$

The minimal surface corresponding to the metric (2.3), in the limit $T \to \infty$ depends only on the x coordinate,

$$X_c = (t, x, \mathbf{0}, Y_c(x)), \tag{2.8}$$

and is qualitatively depicted in Fig.(1). Since the warp factor $F(Y)$ has a minima at $Y = Y^*$, classically the string stays at $Y = Y^*$ except at the end points where it dips to connect to the quark and the antiquark which are located at $Y = 0$. The region of transition is marked d in Fig.(1). For the confining geometries d is expected to grow with L but with $d/L \to 0$ as $L \to \infty$ [24, 25], where one can define d, for example as, that distance for which

$$\frac{Y^* - Y_c(d)}{Y^*} = 10^{-3}. \tag{2.9}$$

In calculating (2.2) using the large D expansion, to start with, we will neglect the region d and consider the fluctuation around the surface given by

$$X_c = (t, x, \mathbf{0}, Y^*), \tag{2.10}$$

(see Fig. (2).) In section 4 we will return to this approximation and make an estimate of the errors



Fig. 1: The classical string configuration in a confining geometry



Fig. 2: Approximation to the classical string configuration used in the large D calculation

introduced by it.

With in this approximation the fluctuations about the minimal surface are given by

$$X = (t, x, \mathbf{X}(t, x), Y^* + \phi(t, x)) \,, \tag{2.11}$$

with $\phi(x, t)$ vanishing at $x = 0$ and at $x = L$. The Nambu-Goto action for such a surface is

$$S_{NG} = T_0 \int_{-T/2}^{T/2} dt \int_0^L dx \, F(Y^* + \phi) \sqrt{\det \left[\gamma_{ab}\right]}, \tag{2.12}$$

where T_0 is the bare string tension, and the induced world-sheet metric γ_{ab} (apart from the warp factor) is

$$\gamma_{ab} = \delta_{ab} + \partial_a \mathbf{X} \cdot \partial_b \mathbf{X} + \partial_a \phi \partial_b \phi. \tag{2.13}$$

It will be convenient to write this metric in terms of two world-sheet matrices,

$$\boldsymbol{g} := g_{ab} = \delta_{ab} + \partial_a \mathbf{X} \cdot \partial_b \mathbf{X}, \tag{2.14}$$

and

$$\boldsymbol{f} := f_{ab} = \partial_a \phi \partial_b \phi. \tag{2.15}$$

Then the action for the world-sheet takes the form

$$S_{NG} = T_0 \int d^2\sigma \, F\left(Y^* + \phi\right) \sqrt{\det g_{ab}} \sqrt{\det\left[\delta_{ab} + \left(g^{-1} f\right)_{ab}\right]}. \tag{2.16}$$

One can think of $T_0 F(Y^* + \phi)$ as a position dependent string tension. Small fluctuation in the Y direction, $\phi(t,x) \neq 0$, increase the string tension. These fluctuations are suppressed compared to the fluctuations $\vec{X} \neq 0, \phi = 0$, for the latter fluctuations keep the string tension fixed at its minimum value. This suggests the following approximate way of evaluating (2.2): Keep transverse fluctuations $\vec{X}$ to all orders but keep only the quadratic fluctuations in ϕ. In this approximation the Nambu-Goto action (2.12) takes the following form

$$S_{NG5} = \frac{1}{l_s^2} \int d^2\sigma \sqrt{\det g_{ab}} + \frac{1}{l_s^2} \int d^2\sigma \sqrt{\det g_{ab}} \left(\frac{1}{2} \mathrm{Tr} \mathbf{g^{-1} f} + \frac{1}{2} M^2 \phi^2\right) \tag{2.17}$$

where M^2 is defined via,

$$F\left(Y^* + \phi\right) = F\left(Y^*\right)\left(1 + \frac{1}{2} M^2 \phi^2\right) + \mathcal{O}\left(\phi^3\right), \tag{2.18}$$

and the fundamental length scale for the problem, l_s, is given by

$$\frac{1}{l_s^2} = T_0 F(Y^*). \tag{2.19}$$

In this approximation, and according to our fundamental assumption (2.2), the expectation value of a Wilson-loop is given by

$$< W[\Gamma] >_{YM} = \int [d\mathbf{X}] \exp\left\{-\frac{1}{l_s^2} \int d^2\sigma \sqrt{\det g_{ab}}\right\} \mathbf{C}\left[g_{ab}\right], \tag{2.20}$$

where

$$\mathbf{C}\left[g_{ab}\right] = \int [d\phi] \exp\left\{-\frac{1}{l_s^2} \int d^2\sigma \sqrt{\det g_{ab}} \left(\frac{1}{2} \mathrm{Tr} \mathbf{g^{-1} f} + \frac{1}{2} M^2 \phi^2\right)\right\}. \tag{2.21}$$

Since the heavy-quark potential, $V\left[L\right]$, is obtained from the expectation value of a rectangular Wilson-Loop

$$\lim_{T\to\infty} < W[\Box_{T\times L}] >_{YM}= \mathcal{N}\exp\left\{-TV\left[L\right]\right\}, \tag{2.22}$$

we see that $\mathbf{C}[g_{ab}]$ contains the contribution to the heavy-quark potential due to the fluctuation of the fundamental string in the fifth dimension. This can be heuristically interpreted as the contribution to the heavy-quark potential due to the fluctuation of the intrinsic thickness of the QCD flux-tube.

Numerical calculations of the ground-state energy of a QCD flux-tube, in lattice gauge theory, matches very well with the ground state energy of the Nambu-Goto string in four euclidean dimensions as given by the Arvis's formula. This suggests that the fluctuations of the string in the fifth dimension should only provide a small correction to the ground-state energy given by the Nambu-Goto string in the four flat dimensions. Therefore, setting $C[\boldsymbol{g}]=$ constant in (2.20), which corresponds to describing the QCD flux-tube by a Nambu-Goto string in four Euclidean dimensions and ignoring the fluctuations in the intrinsic thickness, is a good approximation.

The above observations can be made more precise by evaluating (2.20) in the large D expansion, where as we have noted above D is the number of transverse dimension of the $D+2$ dimensional boundary. As we will see, in the large D expansion of (2.20) the smallness of the correction due to $C[\boldsymbol{g}]$ will be the consequence of the fact that there is only one massive ϕ field while there are D massless transverse fields, X_i.

The large D expansion will be implemented in the standard manner [26], we start with the path-integral

$$Z_5=\int\left[d\mathbf{X}\right]\exp\left\{-\frac{1}{l_s^2}\int d^2\sigma\sqrt{\det g_{ab}}\right\}\mathbf{C}\left[g_{ab}\right], \tag{2.23}$$

and introduce $\boldsymbol{g}$ as an independent degree of freedom in the path integral but with the constraint (2.14). Next we implement the constraint using the Lagrangian multiplier field N. This converts the integration over $\mathbf{X}$ fields into Gaussian integrals and one obtains

$$Z_5=\mathcal{N}_1\int\left[d\boldsymbol{g}\right]\left[d\boldsymbol{N}\right]\exp\left\{-S_{eff}\left[\boldsymbol{g},\boldsymbol{N}\right]\right\}\mathbf{C}[\boldsymbol{g}], \tag{2.24}$$

where

$$S_{eff}[\mathbf{g},\mathbf{N}]=\frac{1}{l_s^2}\int d^2\sigma\left\{\sqrt{\det\mathbf{g}}+\frac{1}{2}N^{ab}(\delta_{ab}-g_{ab})\right\}+\frac{D}{2}\mathrm{Tr}\left[\log\left(-\partial_a N^{ab}\partial_b\right)\right], \tag{2.25}$$

and $\mathcal{N}_1$ is a normalization constant. Further, we can integrate over the ϕ field in (2.21) and write

$$\mathbf{C}[\boldsymbol{g}] = \mathcal{N}_2 \exp\left\{-\frac{1}{2}\mathrm{Tr}\left[\log\left(\hat{\mathbf{A}}\right)\right]\right\}, \tag{2.26}$$

where

$$\hat{\mathbf{A}}\left(\boldsymbol{g}\right) = -\partial_a\sqrt{\det\boldsymbol{g}}g^{-1}_{ab}\partial_b + \sqrt{\det\boldsymbol{g}}M^2, \tag{2.27}$$

and $\mathcal{N}_2$ is another normalization constant.

We would like to evaluate (2.23) in the limit D $\to \infty$ while keeping Dl_s^2 constant. In other words we are interested in the limit

$$\mathrm{D} \to \infty; \quad \bar{l}_s^2 = \text{constant}, \tag{2.28}$$

where $\bar{l}_s^2$ is defined as

$$\bar{l}_s^2 = Dl_s^2. \tag{2.29}$$

The path-integral (2.23) now can be written in the following form

$$Z_5 = \mathcal{N}\int\left[d\boldsymbol{g}\right]\left[d\boldsymbol{N}\right]\exp\left\{-D\,S\left[\boldsymbol{g},\boldsymbol{N}\right]\right\}, \tag{2.30}$$

where $S\left[\boldsymbol{g},\boldsymbol{N}\right]$ has two parts,

$$S\left[\boldsymbol{g},\boldsymbol{N}\right] = \left(S_1\left[\boldsymbol{g},\boldsymbol{N}\right] + \frac{1}{D}S_2\left[\boldsymbol{g}\right]\right), \tag{2.31}$$

with

$$S_1\left[\boldsymbol{g},\boldsymbol{N}\right] = \frac{1}{\bar{l}_s^2}\int d^2\sigma\left\{\sqrt{\det\mathbf{g}} + \frac{1}{2}N^{ab}(\delta_{ab} - g_{ab})\right\} + \frac{1}{2}\mathrm{Tr}\left[\log\left(-\partial_a N^{ab}\partial_b\right)\right], \tag{2.32}$$

and

$$S_2\left[\boldsymbol{g}\right] = \frac{1}{2}\mathrm{Tr}\left[\log\left(\hat{\mathbf{A}}\left(\boldsymbol{g}\right)\right)\right]. \tag{2.33}$$

Therefore in the large D limit, the contribution of the fluctuations of the ϕ field is suppressed by a factor of $1/D$ as compared to the contributions of the fluctuations of the transverse coordinates X_i. From the point of view of the large D expansion, the success of the four-dimensional Nambu-Goto string in reproducing the heavy-quark potential is due to the fact that the corrections to it are suppressed by a factor of $1/D$.

The leading term in the large D expansion of (2.30) is then given by

$$Z_5 = \mathcal{N}_5 \exp\left\{-D\, S_1\left[\bar{\boldsymbol{g}}, \bar{\boldsymbol{N}}\right]\right\}, \tag{2.34}$$

where the $\bar{\boldsymbol{g}}$ and $\bar{\boldsymbol{N}}$ are the fields that minimize the action S_1 alone,

$$\frac{\delta S_1}{\delta \boldsymbol{g}} = 0, \quad \frac{\delta S_1}{\delta \boldsymbol{N}} = 0. \tag{2.35}$$

Minimizing the action S_1 alone is noting but the large D limit of the Nambu-Goto string in a flat Euclidean space [26] and the required solutions are:

$$\bar{\boldsymbol{g}} = \begin{bmatrix} g_1 & 0 \\ 0 & g_2 \end{bmatrix} = \begin{bmatrix} \frac{1-\lambda}{1-2\lambda} & 0 \\ 0 & 1-\lambda \end{bmatrix}, \tag{2.36}$$

$$\bar{\boldsymbol{N}} = \begin{bmatrix} N_1 & 0 \\ 0 & N_2 \end{bmatrix} = \begin{bmatrix} \sqrt{1-2\lambda} & 0 \\ 0 & \frac{1}{\sqrt{1-2\lambda}} \end{bmatrix}, \tag{2.37}$$

where λ depends only on the distance between the quark and the antiquark, L, and is given by

$$\lambda\left(L\right) = \frac{\pi}{24}\frac{\bar{l}_s^2}{L^2} = \frac{\pi D}{24}\frac{l_s^2}{L^2}. \tag{2.38}$$

A fact that will be important for us latter, when we calculate the $1/D$ corrections, the metric $\bar{\boldsymbol{g}}$ has discontinuity at the boundaries, $\sigma = x = 0$ and $\sigma = x = L$, which are the world-lines of the quark and the antiquark (see appendix of [26].) The metric at the boundary is given by

$$\bar{\boldsymbol{g}}_b = \begin{bmatrix} g_{b1} & 0 \\ 0 & g_{b2} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1-2\lambda \end{bmatrix}, \tag{2.39}$$

but interestingly N is continuous,

$$\bar{\boldsymbol{N}} = \begin{bmatrix} \sqrt{\frac{g_{b2}}{g_{b1}}} & 0 \\ 0 & \sqrt{\frac{g_{b1}}{g_{b2}}} \end{bmatrix} = \begin{bmatrix} \sqrt{1-2\lambda} & 0 \\ 0 & \frac{1}{\sqrt{1-2\lambda}} \end{bmatrix}. \tag{2.40}$$

Using the above results, one readily obtains the ground state energy of a Nambu-Goto string in the large D limit [26], which turn out to be the Arvis's formula for the ground state energy of a Nambu-Goto string [4],

$$DS_1\left[\bar{\boldsymbol{g}}, \bar{\boldsymbol{N}}\right] = TV_A\left[L\right], \tag{2.41}$$

$$V_A\left[L\right]=\frac{L}{l_s^2}\sqrt{1-2\lambda\left(L\right)}. \tag{2.42}$$

The large D expansion bears a close resemblance to the large N expansion of multi-component quantum field theories (see for e.g. [27] for a review.) It is also similar in spirit to the mean field approximation (for a mean field analysis of the Nambu-Goto string see [28].) Using the analogy with the large N expansion further, one can ask that is there an analog of the master-field for the $SU(N)$ gauge theories, that is, in the large D limit can the partition function for the Nambu-Goto string written not in terms of sum over all surfaces, weighted by the Nambu-Goto action, but by a single surface. In fact there is such a fictitious mathematical surface, whose metric is given by the boundary metric $\bar{\mathbf{g}}_b$, (2.39),

$$\begin{aligned} Z_{NG} &= \int[d\mathbf{X}]\exp\left\{-\frac{1}{l_s^2}\int d^2\sigma\sqrt{\det g_{ab}}\right\}=\mathcal{N}_{NG}\exp\left\{-\frac{1}{l_s^2}\int_{-T/2}^{T/2}dt\int_0^L dx\sqrt{\det\bar{\mathbf{g}}_b}\right\}, \\ &= \mathcal{N}_{NG}\exp\left\{-\frac{TL}{l_s^2}\sqrt{1-2\lambda(L)}\right\}. \end{aligned} \tag{2.43}$$

This observation will be of great use to us in evaluating the $1/D$ corrections in the next section. The metric $\bar{\mathbf{g}}_b$ should not be thought of as an induced metric, but as an independent metric defining the "master-surface" but which has the right boundary condition, in that when the surface meets the world-line of the quark and the anti-quark then $\bar{g}_{1b}=1=g_{\tau\tau}$ which is the induced metric on the quark and anti-quark world-lines (apart from the overall warp factor $F(Y^*)$ which is absorbed in our definition of l_s (2.19).)

3 The $1/D$ Corrections from Heavy Modes

We will evaluate the $1/D$ corrections arising from the oscillation of the string along the fifth dimension using the observation made in the last section, namely that in the large D limit the partition function of the Nambu-Goto string is given by a single "master surface" with metric given by (2.39).

The partition function of the fundamental string in five dimension, (2.23), can be written as

$$Z_5=Z_{NG}<\mathbf{C}[\boldsymbol{g}]>_{NG}=\mathcal{N}_2 Z_{NG}<\exp\left\{-\frac{1}{2}\mathrm{Tr}\left[\log\left(\hat{\mathbf{A}}[\mathbf{g}]\right)\right]\right\}>_{NG}. \tag{3.1}$$

In the large D limit, this average in turn can be evaluated as

$$Z_5 = \mathcal{N}_5 \exp\left\{-\frac{TL}{l_s^2}\sqrt{1-2\lambda\left(L\right)}\right\} \exp\left\{-\frac{1}{2}\mathrm{Tr}\left[\log\left(\hat{\mathbf{A}}[\bar{\mathbf{g}}_b]\right)\right]\right\}, \tag{3.2}$$

where, using (2.27) and (2.39),

$$\hat{\mathbf{A}}\left[\bar{\boldsymbol{g}}_b\right] = -\sqrt{\frac{g_{b2}}{g_{b1}}}\partial_1^2 - \sqrt{\frac{g_{b1}}{g_{b2}}}\partial_2^2 + \sqrt{\det \bar{\boldsymbol{g}}_b}M^2 \tag{3.3}$$

The trace can be calculated using zeta function regularization in a standard manner (see for e.g [29])

$$-\frac{1}{2}\mathrm{Tr}\left[\log\left(\hat{\mathbf{A}}[\bar{\mathbf{g}}_b]\right)\right] = -TV_M\left(L\right), \tag{3.4}$$

where

$$V_M(L) = \left(-\frac{1}{4}g_{b1}^{1/4}M + \frac{1}{4\pi}\sqrt{\frac{g_{b1}}{g_{b2}}}\int_0^\infty d\omega \log\left(1-\mathrm{e}^{-2L\sqrt{\omega^2+\omega_0^2}}\right)\right), \tag{3.5}$$

and

$$\omega_0 = g_{b2}^{1/2}M. \tag{3.6}$$

Using (2.39) we get

$$V_M\left(L\right) = -\frac{1}{4}M + V_h(L), \tag{3.7}$$

where

$$V_h(L) = \frac{1}{4\pi}\frac{1}{\sqrt{1-2\lambda}}\int_0^\infty d\omega \log\left(1-\mathrm{e}^{-2L\sqrt{\omega^2+(1-2\lambda)M^2}}\right). \tag{3.8}$$

For $LM >> 1$, the integral leads to an exponential decaying term, while for $LM << 1$ one would obtain an additional $1/L$ correction to the heavy quark potential. Such a term is of course precluded both by theoretical arguments [30] and numerical results. In any case for $LM << 1$ our large D analysis fails for then the approximation of replacing $F[Y_c]$ by $F[Y^*]$ in the Nambu-Goto action is no longer valid, equivalently the approximation $d = 0$ in Fig. (1) is no longer valid. Therefore the only region where the above correction is of significance is when $LM \sim 1$, which we can evaluate numerically. In Fig.(3) we show the contribution of difference between $V_A(L)$ and $V_h(L)$ as compared to the Arvis's potential $V_A(L)$ for representative values of M in the units of inverse fermi using $l_s = 0.4$ fermi. Therefore the main qualitative contribution of the heavy modes is to modify the Arvis's potential by an addition of a constant term, $-\frac{1}{4}M$.



Fig. 3: Effect of massive mode on the heavy quark potential, where $\Delta V_{mass} \equiv |V_A - V_h|$

4 The Boundary Effects

Our large D calculation has been done under the approximation where the behavior of the string near the quark and the antiquark was ignored and effectively we have assumed that $d = 0$ in Fig.(1). As a result of this approximation in evaluating (2.2) the fluctuations in the string worldsheets are incorrectly weighted near the vicinity of the quark and the anti-quark. If we relax the approximation $d = 0$ (see Fig.(1) and Fig.(2)), then the minimal surface (2.8) is given by

$$X_c = \Big(t, x, \vec{\mathbf{0}}, Y_c\left(x\right)\Big)\,,$$

and the fluctuations around it are now given by

$$X = \left(t, x, \mathbf{X}\left(t, x\right), Y_c(x) + \phi\left(t, x\right)\right). \tag{4.1}$$

Again, working to the quadratic order in $\phi\left(t, x\right)$, the action for these fluctuations is

$$S = T_0 \int_{-T/2}^{T/2} dt \int_0^L dx F\left[Y_c\left(x\right)\right] \left(1 + \frac{1}{2} g_{22}^{-1}\left(\partial_x Y_c\right)\right) \sqrt{\det \mathbf{g}} + S_h[\mathbf{g}, \phi], \tag{4.2}$$

where $S_h[\mathbf{g}, \phi]$ is a quadratic functional of $\phi(t, x)$ whose explicit form is given in the Appendix (A.) Since the classical string configuration, $Y_c(x)$, deviates from $Y = Y^*$ only near the boundary, therefore we can write the warp factor as

$$F\left[Y_c\right] = F^* + F^*\eta\left[Y_c\right], \tag{4.3}$$

where, to a very good approximation, $\eta\left[Y_c\right]$ has support only in the regions $0 < x < d$ and $L - d < x < L$. The action then can be written as

$$\begin{aligned} S\left[\mathbf{X}, \phi; F\left(Y\right)\right] = \frac{1}{l_s^2}\int_{-T/2}^{T/2} dt \int_0^L dx \sqrt{\det \mathbf{g}} + \\ \frac{1}{l_s^2}\int_{-T/2}^{T/2} dt \int_0^d dx \mathbf{B}\left[Y_c(x), \mathbf{g}\right]\sqrt{\det \mathbf{g}} + \frac{1}{l_s^2}\int_{-T/2}^{T/2} dt \int_{L-d}^L dx \mathbf{B}\left[Y_c(x), \mathbf{g}\right]\sqrt{\det \mathbf{g}} + \\ S_h\left[\mathbf{g}, \phi\right], \end{aligned} \tag{4.4}$$

the world-sheet matrix $\mathbf{g}$ is given by (2.14) while $\mathbf{B}\left[Y_c\left(x\right), \mathbf{g}\right]$ and $S_h\left[\mathbf{g}, \phi\right]$ are given by (A.4) and (A.5) respectively. One can repeat the analysis of section (2), and again one finds that the contribution of the fluctuations $\phi\left(t, x\right)$ along the fifth dimension, which are governed by $S_h\left[\mathbf{g}, \phi\right]$, are suppress by a factor of $1/D$ as compared to the mass-less fluctuations, $\mathbf{X}\left(t, x\right)$, which are along the directions transverse to the QCD flux-tube. The dynamics of these fluctuations in the large D limit is governed by

$$\begin{aligned} S\left[\mathbf{X}; F\left[Y_c\left(x\right)\right]\right] = \frac{1}{l_s^2}\int_{-T/2}^{T/2} dt \int_0^L dx \sqrt{\det \mathbf{g}} + \\ \frac{1}{l_s^2}\int_{-T/2}^{T/2} dt \int_0^d dx \mathbf{B}\left[Y_c(x), \mathbf{g}\right]\sqrt{\det \mathbf{g}} + \frac{1}{l_s^2}\int_{-T/2}^{T/2} dt \int_{L-d}^L dx \mathbf{B}\left[Y_c(x), \mathbf{g}\right]\sqrt{\det \mathbf{g}}. \end{aligned} \tag{4.5}$$

Restricting to the cases for which $d/L << 1$, and assuming that $\mathbf{B}\left[Y_c(x), \mathbf{g}\right]$ has a finite value at the boundary, we can approximate the last two terms by expanding the integrand about its boundary value to obtain

$$\begin{aligned} S\left[\mathbf{X}; F\left[Y_c\left(x\right)\right]\right] = \frac{1}{l_s^2}\int_{-T/2}^{T/2} dt \int_0^L dx \sqrt{\det \mathbf{g}} + \\ \frac{d}{l_s^2}\int_{-T/2}^{T/2} dt \left(\mathbf{B}\sqrt{\det \mathbf{g}}\right)_{x=0} + \frac{d}{l_s^2}\int_{-T/2}^{T/2} dt \left(\mathbf{B}\sqrt{\det \mathbf{g}}\right)_{x=L} + \cdots, \end{aligned} \tag{4.6}$$

where dots represent the terms coming from derivatives of the integrand. From the point of view of the boundary theory, this is an effective action for a QCD flux-tube, and it is nothing but the

Nambu-Goto string in four flat dimensions but supplemented with boundary terms.

Let us now come back to our assumption about $\mathbf{B}\left[Y_c(x),\mathbf{g}\right]$ and inquire if it is really finite at the boundary[3]. From (A.4) we have

$$B[Y_c(x),\mathbf{g}] = \left(\frac{1}{2}g_{22}^{-1}\partial_x Y_c + \eta\left(x\right)(1+\frac{1}{2}g_{22}^{-1}\partial_x Y_c)\right), \tag{4.7}$$

if the behavior of the warp factor $F(Y)$ near the boundary is similar to that for the AdS_5 space then $\eta(x)$ near the boundary behaves as

$$\eta\left(x\right) \sim \frac{1}{Y_c\left(x\right)^2} \sim \frac{1}{x^2}, \tag{4.8}$$

where we have focused on the behavior of the fundamental string near the quark situated at the origin. This divergence in $\eta\left(x\right)$ is known to be related to the divergent self-energy of the quark and the anti-quark on which the fundamental string terminates [23]. Therefore it is plausible that after implementing the rernomalization procedure that removes this divergence we should be left with a finite boundary term of the form

$$B[Y_c(x),\mathbf{g}]_{boundary} = Z_{finite}\left(g_{22}^{-1}\partial_x Y_c\right)_{boundary} = \mathbf{C}\left(g_{22}^{-1}\right)_{boundary}, \tag{4.9}$$

where $\mathbf{C}$ is an unknown constant.

If we assume that this finite boundary term can be treated as a small perturbation to the leading term, the Nambu-Goto term, in (4.6), then its effect can be estimated in the large D limit. For then the partition function in the large D limit takes the form

$$\begin{aligned} Z_d &\equiv \int\left[d\mathbf{X}\right]\exp\left\{-S\left[\mathbf{X};F\left[Y_c\left(x\right)\right]\right]\right\} \\ &= Z_{NG}\left\langle\exp\left(-\frac{d}{l_s^2}\int_{-T/2}^{T/2}dt\left(B\sqrt{\det\mathbf{g}}\right)_{x=0} - \frac{d}{l_s^2}\int_{-T/2}^{T/2}dt\left(\mathbf{B}\sqrt{\det\mathbf{g}}\right)_{x=L}\right)\right\rangle_{Z_{NG}}. \end{aligned} \tag{4.10}$$

Evaluating it in the large D limit, using (2.43),

$$Z_d = \mathcal{N}_d\exp\left\{-\frac{TL}{l_s^2}\sqrt{\bar{\mathbf{g}}_b} - \frac{d}{l_s^2}T\sqrt{\bar{\mathbf{g}}_b}\mathbf{B}\left(0;\bar{\mathbf{g}}_b\right) - \frac{d}{l_s^2}T\sqrt{\bar{\mathbf{g}}_b}\mathbf{B}\left(L,\bar{\mathbf{g}}_b\right)\right\}, \tag{4.11}$$

---

[3] I would like to thank Ofer Aharony for raising this question.



Fig. 4: Fluctuation leading to the formation of the caps of the flux-tube

which leads to the following static potential

$$V_0[L] = \frac{L}{l_s^2}\left(1 + \frac{d}{L}\frac{2\mathbf{C}}{\bar{g}_{b2}}\right)\sqrt{\det \bar{\mathbf{g}}_b}. \tag{4.12}$$

In a confining geometry it is expected that d/L vanishes as $L \to \infty$ [24, 25], with this in mind let us assume that

$$\frac{d}{L} \sim \left(\frac{l_s}{L}\right)^{\alpha}, \tag{4.13}$$

where α is an unknown positive real number. Then the potential can be written as

$$V_0\left[L\right] = \frac{L}{l_s^2}\left(1 + \beta\left(\frac{l_s}{L}\right)^{\alpha}\frac{1}{1-2\lambda\left(L\right)}\right)\sqrt{1-2\lambda\left(L\right)}, \tag{4.14}$$

where α and β are two unknown parameter which depends on the details of the confining geometry and $\lambda(L)$ is given by (2.38.) Since we haves assumed that the boundary terms are a small corrections to the Nambu-Goto term in (4.6) therefore we expect $\beta << 1$. Heuristically speaking, the effect of the boundary terms is to give the string tension of the QCD flux-tube a very mild dependence on the length of the flux tube.

The need for boundary terms in a QCD flux-tube is also suggested when one considers the possible interaction of the QCD flux-tube with the spin of the quark and the anti-quark [31]. If one tries to model this interaction in an effective string description of a QCD flux-tube then one is forced to introduce a boundary term [32] which takes into account the intrinsic thickness of the QCD flux-tube.

Finally let us consider the fluctuations depicted in Fig. (4), these are the fluctuations of the string in the extra dimension that have not been included in the large D calculation. In fact these fluctuations cannot be described in the static gauge we have used. Though the following qualitative effects of these fluctuations is suggested by the idea that the holographic projection of a fundamental string on to the boundary is a thick flux-tube whose thickness depends on the distance of the fundamental string from the boundary [15]. This then implies that the shape of the flux-tube corresponding to the classical configuration of the fundamental string is the one shown in Fig. (1). The shape of the flux-tube in Fig. (2) is very different from expected shape of a QCD flux-tube as shown in Fig. (4), this suggests that the effect of these fluctuations in the region $0 < x < d$ and in the region $L - d < x < L$ is to make the flux in the flux-tube penetrate the region of the order of ξ beyond the quark and the anti-quark. ξ being the correlation length of the gauge field, or the inverse glue-ball mass, which one expects to be of the order of magnitude of l_s.

5 Summary and Conclusions

The key features of the heavy quark potential as suggested by the gauge/gravity duality in the large D expansion can be summarized from (3.7,4.14),

$$V[L] = \left((1 + f(L)) \frac{L}{l_s^2} \sqrt{\left(1 - \frac{2\pi}{12} \frac{l_s^2}{L^2}\right)} \right) + \left(-\frac{1}{4} M + V_h[L] \right), \tag{5.1}$$

where from (4.14) we see that the leading behavior of $f(L) \approx 1 + \beta \left(\frac{l_s}{L}\right)^{\alpha} + \cdots$. The contribution from the heavy modes is contained in $V_h[L]$ which is given by (3.8) and is shown in Fig. (3.) We have grouped the leading corrections in the large D expansion in the first parenthesis while the second parenthesis contains the $1/D$ corrections arising from the oscillations of the fundamental string in the fifth dimension. As we have noted in the introduction that there are possible $1/D$ and higher corrections arising from the massless modes in non-critical dimensions, which we have ignored in our analysis. There analysis and to understand the manner in which they vanish for the critical dimension is an important and an interesting problem to which we hope to come in the future.

There are two key qualitative features of $V[L]$. Firstly, the correction to the Arvis's potential by the factor $f(L)$. Though in our investigation we cannot fix the precise form of $f(L)$ we can see clearly the reason for its existence. It arises from the simultaneous requirement that the fundamental string wants to lie at a fixed value $Y = Y^*$, to minimize its energy, and the requirement that the fundamental string has to terminate on the boundary, $Y = 0$, where the quark and the anti-quark are placed. Thus, the origin of $f(L)$ is in the behavior of the flux-tube in the vicinity of the quark and the anti-quark. Thought the present numerical results are consistent with $f(L) = 0$, it will be very interesting to fit the future numerical calculation of the heavy quark potential to the above form for the heavy quark potential. The other interesting feature of (5.1) is the existence of the constant term in the potential, $-M/4$. At the first sight the existence of constant term in potential is of no consequence, but as has been emphasized in [33, 34], that once the renormalization prescription for the expectation value of the Wilson loop is appropriately fixed, the constant arises only from the corrections to the Nambu-Goto string[4]. It is worth recalling that there is no constant term in the ground state energy of a Nambu-Goto string as given by the Arvis's formula.

The gauge/gravity duality provides us with a very useful geometrical way of understanding the confining property of QCD. Even though this description is largely heuristic, as we lack the gravitational dual of QCD that includes both confinement and the asymptotic freedom, still it gives us various new insight into the formation and the dynamics of the QCD flux-tube. These insights follow from the geometrical idea that the QCD flux-tube is a holographic projection of a fundamental five-dimensional string which terminates on the quark and the anti-quark that live on the four-dimensional boundary of a curved five-dimensional space. For example, it explains the success of the Nambu-Goto description of the QCD flux-tube, even though the flux-tube is not a one-dimensional fundamental string. From the geometrical point of view, the fluctuations of intrinsic thickness of the flux-tube, which distinguishes it from a fundamental string, corresponds to fluctuations of the fundamental string along the fifth-dimension and these fluctuations are massive and hence suppressed. Though the explanation again is heuristic, for the mass of these modes cannot be predicated and one has to appeal to dimension analysis and argue that they must be of the order of the glueball mass. Further the same geometric picture suggests that the Nambu-Goto description of the QCD flux-tube has to be supplemented by boundary terms, for in the immediate vicinity of the quark and the anti-quark the fluctuations of the fundamental string is qualitatively different from that of a Nambu-Goto string in flat four dimensional space. It is interesting to note that these boundary terms appear not because of integrating out the heavy mode, corresponding to the oscillations in fifth dimension, but because of the behavior of the massless modes near the boundary. Better understanding of these boundary terms therefore may also give us some clue as to how does gauge/gravity duality implements the asymptotic freedom.

Acknowledgments

I would like to thank Ofer Aharony for his very useful comments to the preliminary version of this paper. I have also greatly benefited from my discussions with the participants of QCD-String

---

[4] I would like to thank R. D. Pisarski for pointing this out to me.

workshop, Trento, 2010, and I would like to thank them all.

A Appendix: Fluctuations Around the Minimal Surface

We want to consider the fluctuations (4.2) around the minimal surface (2.8), their action is given by

$$S = \int_{-T/2}^{T/2} dt \int_0^L dx F\left[Y_c(x) + \phi(x,t)\right] \sqrt{\det\left(\mathbf{g} + \mathbf{f}_c\right)}, \tag{A.1}$$

where $\mathbf{g}$ is given by (2.14), while

$$(\mathbf{f}_c)_{ab} = \partial_a \left(Y_c\left(x\right) + \phi\left(t,x\right)\right) \partial_b \left(Y_c\left(x\right) + \phi\left(t,x\right)\right). \tag{A.2}$$

If one expands the action to second order in $\phi\left(t,x\right)$ and use the equations of motion for $Y_c(x)$ then one obtains

$$S\left[\mathbf{X},\phi;F\left(Y\right)\right] = \frac{1}{l_s^2}\int_{-T/2}^{T/2} dt \int_0^L dx \sqrt{\det\mathbf{g}} + \frac{1}{l_s^2}\int_{-T/2}^{T/2} dt \int_0^L dx \mathbf{B}\left[Y_c(x),\mathbf{g}\right]\sqrt{\det\mathbf{g}} + S_h\left[\mathbf{g},\phi\right], \tag{A.3}$$

where

$$B[Y_c(x),\mathbf{g}] = \left(\frac{1}{2}g_{22}^{-1}\partial_x Y_c + \eta\left(x\right) + \frac{1}{2}\eta(x) g_{22}^{-1}\partial_x Y_c\right), \tag{A.4}$$

and $\eta(x)$ is given by (4.3) while

$$S_h\left[\mathbf{g},\phi\right] = \int_{-T/2}^{T/2} dt \int_0^L dx \frac{1}{2}\left(A\left[\mathbf{g},\partial_x Y_c\right]\left(\partial_t\phi\right)^2 + B\left[\mathbf{g},\partial_x Y_c\right]\left(\partial_x\phi\right)^2 + \right) + \int_{-T/2}^{T/2} dt \int_0^L dx \frac{1}{2}\left(C\left[\mathbf{g},\partial_x Y_c\right]\left(\partial_t\phi\right)\left(\partial_x\phi\right) + M^2\left[\mathbf{g},\partial_x Y_c\right]\phi^2\right). \tag{A.5}$$

The factors A, B, C and M^2 are calculable functionals of $\mathbf{g}$ and $\partial_x Y$ but we will not require their explicit form.

References

[1] N. D. Hari Dass and P. Majumdar, *String-like behaviour of 4d SU(3) Yang-Mills flux tubes*, *JHEP* **10** (2006) 020, [hep-lat/0608024].

[2] B. B. Brandt and P. Majumdar, *Spectrum of the QCD flux tube in 3d SU(2) lattice gauge theory*, *Phys. Lett.* **B682** (2009) 253, [arXiv:0905.4195].

[3] B. B. Brandt, *Probing boundary-corrections to Nambu-Goto open string energy levels in 3d SU(2) gauge theory*, *JHEP* **1102** (2011) 040, [arXiv:1010.3625].

[4] J. F. Arvis, *The Exact q anti-q Potential In Naumbu String Theory*, *Phys. Lett.* **B127** (1983) 106.

[5] K. G. Wilson, *Quantum chromodynamics on a lattice*, . Presented at Cargese Summer Institute, Cargese, France, Jul 12- 31, 1976.

[6] F. David, *Introduction to the statistical mechanics of random surfaces and membranes*, 1991. Lectures given at 8th Jerusalem Winter School for Theoretical Physics, Two-Dimensional Gravity and Random Surfaces, Jerusalem, Israel, Dec 27, 1990 - Jan 4, 1991.

[7] S. Dubovsky, R. Flauger, and V. Gorbenko, *Effective string theory revisited*, arXiv:1203.1054.

[8] J. Polchinski and A. Strominger, *Effective string theory*, *Phys. Rev. Lett.* **67** (1991) 1681.

[9] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri, and Y. Oz, *Large n field theories, string theory and gravity*, *Phys. Rept.* **323** (2000) 183, [hep-th/9905111].

[10] J. M. Maldacena, *The large N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231, [hep-th/9711200].

[11] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, *Gauge theory correlators from non-critical string theory*, *Phys. Lett.* **B428** (1998) 105–114, [hep-th/9802109].

[12] E. Witten, *Anti-de Sitter space and holography*, *Adv.Theor.Math.Phys.* **2** (1998) 253–291, [hep-th/9802150].

[13] E. Witten, *Anti-de sitter space, thermal phase transition, and confinement in gauge theories*, *Adv. Theor. Math. Phys.* **2** (1998) 505–532, [hep-th/9803131].

[14] U. H. Danielsson, E. Keski-Vakkuri, and M. Kruczenski, *Vacua, Propagators, and Holographic Probes in AdS/CFT*, *JHEP* **01** (1999) 002, [hep-th/9812007].

[15] J. Polchinski and L. Susskind, *String theory and the size of hadrons*, hep-th/0112204.

[16] V. Vyas, *Intrinsic Thickness of QCD Flux-Tubes*, arXiv:1004.2679.

[17] H. B. Nielsen and P. Olesen, *Vortex-line models for dual strings*, *Nucl. Phys.* **B61** (1973) 45.

[18] O. Aharony, *Private communication*, .

[19] O. Aharony and E. Karzbrun, *On the effective action of confining strings*, *JHEP* **06** (2009) 012, [arXiv:0903.1927].

[20] O. Aharony and M. Field, *On the effective theory of long open strings*, *JHEP* **1101** (2011) 065, [arXiv:1008.2636].

[21] K. G. Wilson, *Confinement of quarks*, *Phy. Rev. D* **10** (1974) 2445.

[22] J. M. Maldacena, *Wilson loops in large n field theories*, *Phys. Rev. Lett.* **80** (1998) 4859, [hep-th/9803002].

[23] L. Susskind and E. Witten, *The holographic bound in anti-de Sitter space*, hep-th/9805114.

[24] J. Greensite and P. Olesen, *Worldsheet fluctuations and the heavy quark potential in the AdS/CFT approach*, *JHEP* **04** (1999) 001, [hep-th/9901057].

[25] Y. Kinar, E. Schreiber, J. Sonnenschein, and N. Weiss, *Quantum fluctuations of Wilson loops from string models*, *Nucl. Phys.* **B583** (2000) 76, [hep-th/9911123].

[26] O. Alvarez, *The Static Potential in String Models*, *Phys. Rev.* **D24** (1981) 440.

[27] S. Coleman, *Aspects of Symmetry: Selected Erice Lectures*. Cambridge University Press, Feb., 1988.

[28] Y. Makeenko, *Qcd string as an effective string*, arXiv:1206.0922.

[29] V. V. Nesterenko and I. G. Pirozhenko, *Justification of the zeta function renormalization in rigid string model*, *J. Math. Phys.* **38** (1997) 6265–6280, [hep-th/9703097].

[30] M. Luscher and P. Weisz, *String excitation energies in SU(N) gauge theories beyond the free-string approximation*, *JHEP* **07** (2004) 014, [hep-th/0406205].

[31] J. Kogut and G. Parisi, *Long-range spin-spin forces in gauge theories*, *Phys. Rev. Letts.* **47** (1981) 1089.

[32] V. Vyas, *Spin-String Interaction in QCD Strings*, *Phys. Rev.* **D78** (2008) 045003, [arXiv:0704.2707].

[33] Y. Hidaka and R. D. Pisarski, *Zero Point Energy of Renormalized Wilson Loops*, *Phys.Rev.* **D80** (2009) 074504, [arXiv:0907.4609]. 19 pages, RevTeX.

[34] U. Kol and J. Sonnenschein, *Can holography reproduce the QCD Wilson line?*, *JHEP* **1105** (2011) 111, [arXiv:1012.5974].